

# TopleVo 500

Levofloxacin  
500 mg Tablet

**TopleVo 500** (Levofloxacin) is a synthetic, broad-spectrum, third generation fluoroquinolone antibiotic. Chemically, Levofloxacin is a chiral fluorinated carboxyquinolone.

## COMPOSITION

Each film coated tablet contains Levofloxacin Hemihydrate INN equivalent to Levofloxacin 500 mg.

## INDICATION

**TopleVo 500** (Levofloxacin) is indicated for the treatment of respiratory tract infections including acute bacterial sinusitis, acute exacerbations of chronic bronchitis, community-acquired pneumonia and nosocomial pneumonia. It is also used in the treatment of uncomplicated and complicated skin & soft tissue infections, chronic bacterial prostatitis and uncomplicated & complicated urinary tract infections (mild to moderate). It is indicated for the treatment of adults (18 years of age).

## DOSAGE AND ADMINISTRATION

*Acute sinusitis* : 500 mg daily for 10-14 days.

*Acute exacerbations of chronic bronchitis* : 250-500 mg daily for 7-10 days.

*Community-acquired pneumonia* : 500 mg once or twice daily for 7-14 days.

*Nosocomial pneumonia* : 750 mg once daily for 7-14 days.

*Skin & soft tissue infections* : 500 mg once or twice daily for 7-14 days.

*Chronic prostatitis* : 500 mg once daily for 28 days.

*Urinary tract infections* : 250 mg daily for 7-10 days (for 3 days in uncomplicated infections).

## SIDE EFFECT

**TopleVo 500** (Levofloxacin) is generally well tolerated. However, a few side effects can usually be seen. Side effects include nausea, vomiting, diarrhea, abdominal pain, flatulence and rare occurrence of phototoxicity (0.1%). Side effects that may be seen very rarely include - tremors, depression, anxiety, confusion etc.

## PRECAUTION

Adequate amount of water should be taken during treatment period to avoid concentrated form of urine. Dose adjustment is required for patients with impaired renal function.

## CONTRAINDICATION

**TopleVo 500** (Levofloxacin) is contraindicated in patients with a history of hypersensitivity to Levofloxacin and other quinolone antimicrobial agents or any other components of this product.

## DRUG INTERACTION

Antacids, iron and adsorbents reduce absorption of Levofloxacin. NSAIDs may increase the risk of CNS stimulation. Warfarin may increase the risk of bleeding.

## USE IN PREGNANCY & LACTATION

US FDA pregnancy category of Levofloxacin is C. So, Levofloxacin should be avoided in pregnancy and lactation unless the potential benefits to the other outweigh the possible risks to the fetus.

## STORAGE CONDITION

Keep in a dry place away from light and heat. Keep out of reach of children.

## HOW SUPPLIED

Each box contains 30 tablets in Alu-PVC blister pack.

Manufactured by:

**NIPRO JMI Pharma Ltd.**  
Comilla, Bangladesh.

