

# Lijenta 5

Linagliptin  
5 mg Tablet

## COMPOSITION

Each film coated tablet contains Linagliptin INN 5 mg.

## INDICATION

Lijenta is indicated in the treatment of type 2 diabetes mellitus to improve glycaemic control in adults.

As *monotherapy*-in patients inadequately controlled by diet and exercise alone and for whom metformin is inappropriate due to intolerance or contraindicated due to renal impairment.

As *combination therapy*-in combination with metformin when diet and exercise plus metformin alone do not provide adequate glycaemic control-in combination with a sulphonylurea and metformin when diet and exercise plus dual therapy with these medicinal products do not provide adequate glycaemic control.

## DOSAGE AND ADMINISTRATION

5 mg once daily. If added to metformin, the dose of metformin should be maintained and linagliptin administered concomitantly. When used in combination with a sulphonylurea, a lower dose of the sulphonylurea may be considered to reduce the risk of hypoglycaemia.

*Patients with renal impairment:* no dose adjustment required. Linagliptin can be taken with or without a meal at any time of the day.

## CONTRAINDICATION

Hypersensitivity to the active substance or to any of the excipients.

## PRECAUTION

Linagliptin should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

## SIDE EFFECT

There may be hypoglycaemia, nasopharyngitis, cough and pancreatitis in combination with metformin and sulfonylurea.

## DRUG INTERACTION

Linagliptin is a weak competitive and a weak to moderate mechanism-based inhibitor of CYP isozyme CYP3A4, but does not inhibit other CYP isozymes. The risk for clinically meaningful interactions by other medicinal products on linagliptin is low and in clinical studies linagliptin had no clinically relevant effect on the pharmacokinetics of metformin, glyburide, simvastatin, warfarin, digoxin or oral contraceptives.

## PREGNANCY AND LACTATION

Avoid use during pregnancy. A risk to the breast-fed child cannot be excluded.

## STORAGE CONDITION

Keep in a dry place away from light and heat. Keep out of the reach of children.

## HOW SUPPLIED

Each box contains 10 tablets in Alu-Alu blister pack.



Manufactured by:

**NIPRO JMI Pharma Ltd.**  
Comilla, Bangladesh.