

Fexten

Fexofenadine Hydrochloride USP

Fexten (Fexofenadine Hydrochloride) is an antihistamine with selective peripheral H₁-receptor antagonist activity. It is rapidly absorbed after oral administration and peak plasma concentration is reached in 2-3 hours. It does not appear to cross the blood brain barrier.

COMPOSITION

Fexten 60 Tablet : Each film coated tablet contains Fexofenadine Hydrochloride USP 60 mg.

Fexten 120 Tablet : Each film coated tablet contains Fexofenadine Hydrochloride USP 120 mg.

Fexten Suspension : Each 5 ml suspension contains Fexofenadine Hydrochloride USP 30 mg.

INDICATION

It is indicated for the relief of symptoms associated with seasonal allergic rhinitis in adults and children. It is also indicated for the treatment of uncomplicated skin manifestations of chronic idiopathic urticaria.

DOSAGE AND ADMINISTRATION

Disease	Age group	Fexten Tablet	Fexten Suspension	In case of impaired renal function
Seasonal Allergic Rhinitis	Adults and children 12 years and older	60 mg twice daily or 120 mg once daily or 180 mg once daily	-----	60 mg once daily
	Children from 6 to 11 years	30 mg twice daily or 60 mg once daily	-----	30 mg once daily
	Children from 2 to 11 years	-----	30 mg or 5 ml (1tsp) twice daily	30 mg or 5 ml (1tsp) once daily
Chronic Idiopathic Urticaria	Adults and children 12 years and older	60 mg twice daily or 120 mg once daily or 180 mg once daily	-----	60 mg once daily
	Children from 6 to 11 years	30 mg twice daily or 60 mg once daily	-----	30 mg once daily
	Children from 6 months to less than 2 years	-----	15 mg or 2.5 ml (1/2 tsp) twice daily	15 mg or 2.5 ml (1/2 tsp) once daily
	Children from 2 to 11 years	-----	30 mg or 5 ml (1tsp) twice daily	30 mg or 5 ml (1tsp) once daily

SIDE EFFECT

Common side effects are headache, fatigue, drowsiness, nausea, dry mouth and gastrointestinal disturbances.

PRECAUTION

Caution should be exercised in elderly patient and patient with decreased renal function.

CONTRAINDICATION

Contraindicated in patients with known hypersensitivity to Fexofenadine Hydrochloride or any of its ingredients.

DRUG INTERACTION

Plasma concentration of Fexofenadine Hydrochloride have been increased when given with erythromycin or ketoconazole. Aluminium and magnesium hydroxide containing antacid reduces the absorption of Fexofenadine Hydrochloride.

USE IN PREGNANCY AND LACTATION

US FDA pregnancy category of Fexofenadine Hydrochloride is C. So, Fexofenadine Hydrochloride should be avoided in pregnancy and lactation unless the potential benefits to the other outweigh the possible risks to the fetus.

STORAGE CONDITION

Keep in a dry place away from light and heat. Keep out of the reach of children.

HOW SUPPLIED

Fexten 60 Tablet : Each box contains 30 tablets in Alu-Alu blister pack.

Fexten 120 Tablet : Each box contains 30 tablets in Alu-Alu blister pack.

Fexten Suspension : Each Amber PET bottle contains 50 ml suspension.