

OxyFast

Nasal Spray

Oxymetazoline
Hydrochloride USP 0.05%

COMPOSITION

Each actuation delivers Oxymetazoline Hydrochloride USP 25 microgram.

PHARMACOLOGY

Oxymetazoline Hydrochloride is an imidazoline derivative sympathomimetic amine. It is a direct agonist at α -adrenoceptors but has no action on β -adrenoceptors. It is used as a topical agent on the nasal mucosa, produces a rapid and long acting vasoconstriction of the arterioles, thus reducing the blood flow and diminishing swelling of the mucosa. This results in improved potency of the airway and better drainage of the nasal sinus.

INDICATION

OxyFast (Oxymetazoline Hydrochloride) is indicated for the treatment of nasal congestion associated with acute or chronic rhinitis, common cold, sinusitis, upper respiratory allergies, epistaxis and hay fever. It is also used as nasal decongestant in otitis media.

DOSAGE & ADMINISTRATION

OxyFast Nasal Spray is recommended for intranasal use only.

Adults and children (6 years of age and older): 2 or 3 sprays in each nostril twice daily for 3-5 days.

Elderly: There is no need for dosage reduction in the elderly for nasal administration.

HOW TO USE

Step 01: Remove the dust cap and neck lock



Step 02: During application, tilt your head slightly forward and keep the container upright then use the nasal spray



Step 03: After each use, gently wipe the spray tip with clean & soft cloth or tissue paper. Finally, set the dust cap in the previous place



CONTRAINDICATION

It is contraindicated in patients with cardiovascular disease, hyperthyroidism, prostatic enlargement, angle closure glaucoma and hypersensitivity to any of its ingredients.

WARNING & PRECAUTION

Keep away from the eyes. The spray should not be used for longer than seven days without medical advice. Frequent or prolonged use may cause nasal congestion to recur or worsen.

SIDE EFFECTS

Occasionally it may cause local irritation and dryness of the mouth and throat. Temporarily discomfort such as burning, stinging, sneezing or increase in nasal discharge may occur.

USE IN PREGNANCY & LACTATION

The safety of Oxymetazoline Hydrochloride Nasal Spray during pregnancy and lactation has not been established, but is not thought to constitute a hazard during those periods.

USE IN ADOLESCENTS AND CHILDREN

It is not suitable for children below 6 years of age.

DRUG INTERACTION

Oxymetazoline Hydrochloride Nasal Spray should not be given to patients being treated with MAO inhibitors or within 14 days of ceasing such treatment.

OVERDOSE

Overdose may give rise to local irritation and rebound congestion. Treatment need only be symptomatic.

STORAGE CONDITION

Keep below 30°C temperature in a dry place, protected from light. Do not refrigerate or freeze. Keep out of the reach of children.

PACKING

Each bottle contains Oxymetazoline Hydrochloride for 200 metered sprays.